

Testudines

Emydoidea blandingii -- Blanding's Turtle

Search for another IL amphibian or reptile


Search for IL amphibians & reptiles by county

Go to the INHS Amphibian & Reptile Collection homepage

Search the INHS collections data for amphibians & reptiles

Search the INHS Type Specimen Data for amphibians & reptiles

Cook County, IL; photo by Mike Redmer



Purple shade indicates vouchered specimens. Light blue (cyan) shade indicates photographic records. Yellow shade indicates verified sighting. Slanted hatch indicates pre-1980 records only

NOTE: Not all specimens upon which these maps are based have been verified.

Key Characters: Bright yellow chin and throat; notched upper jaw; hinged plastron.

Similar species: Box turtles, spotted turtle.

Description: Medium-sized (up to 24 cm CL) turtle with dark shell. Head and carapa profusely decked with light spots and dashes. Each plastral scute patterned with a lar dark blotch usually bordered with yellow but, in older individuals, entire plastron may be black. Pleural scutes of young unpatterned or have faint radiating streaks. Disproportionately longer tail than box turtles. Male with concave plastron, heavier ta with cloacal opening behind edge of carapace, and dark markings on upper mandible

Habitat: Quiet waters in marshes, prairie wetlands, wet sedge meadows, and shallow vegetated portions of lakes.

Natural History: Usually found in and around water, but moves long distances overland. Long-lived, up to 77 years. Chiefly carnivorous, eating snails, insects, crayfish, and vertebrates. Female nests in late May and June, laying one clutch of abo 12 hard-shelled, ellipsoidal eggs (ca. 30 x 20 mm).

Status: State threatened. Major threat has been habitat destruction. Relatively common in appropriate habitat from Illinois River northward. Rare farther south (see distribution map, above).

©Illinois Natural History Survey
Send Questions and Comments to:cbdadmin@inhs.uiuc.edu.
Last updated Wednesday, 07-Apr-2004 10:19:45 CDT